

QUANTUM ROTATIONAL SPECTRA AND CLASSICAL ROTORS

Marek MIŚKIEWICZ*, Andrzej GÓŹDŹ†

*Department of Theoretical Physics,
University of Maria Curie – Skłodowska
Lublin, Poland*

Jerzy DUDEK‡

*Institut de Recherches Subatomiques,
IN2P3-CNRS and Université Louis Pasteur, Strasbourg
France*

We consider the generalized rotor Hamiltonians capable of describing quantum systems invariant with respect to symmetry point-groups that go beyond the usual D_2 -symmetry of a tri-axial rotor. We discuss the canonical de-quantisation procedure to obtain the classical analogs of the original quantum Hamiltonians. Classical and quantum solutions to the Hamiltonians relevant in the nuclear physics applications are illustrated and compared using the 'usual' (D_2) and an 'exotic' (T_d) symmetries.

GENERAL ASPECTS

In this paper we consider the so-called *generalized rotor hamiltonians*, i.e. hamiltonians capable of describing the rotating quantum objects that are invariant under an arbitrary classical point group of symmetry. The word *generalized* implies that we go beyond the properties of the standard D_2 -symmetric ('tri-axial') rotors. We also study the classical analogs of the corresponding quantum objects - 'generalised classical rotors' and the corresponding trajectories - solutions to the related classical Hamilton equations of motion.

Hamiltonians of this kind are of special interest in the recent nuclear physics applications not to mention the traditional ones related to the molecular physics; realistic microscopic

* marekm@kft.umcs.lublin.pl

† gozdz@neuron.umcs.lublin.pl

‡ Jerzy.Dudek@ires.in2p3.fr

calculations predict the possibility that some excited nuclear states may be characterised by the high tetrahedral and/or octahedral symmetries, cf. Refs.[1] and also in: N. Schunck and J. Dudek, these Proceedings.

The point groups are subgroups of the orthogonal group, $O(3)$; for the latter one can define operations of the left- and the right-shifts [2]. Introducing laboratory and intrinsic frames it can be easily shown that the left-shift operation corresponds to the rotation in the laboratory- and the right-shift operation to the one in the body-fixed frame, respectively. One can demonstrate that the generators of the left-shift should be interpreted as the angular momentum operators in the laboratory frame, and the generators of the right-shift as the operators of the angular momentum in the intrinsic frame. These two notions allow to introduce the definitions of the rotor hamiltonians in the mathematically consistent, precise manner. Here we will not go more into any mathematical details leaving the corresponding discussion to a forthcoming publication.

The commutation-relations of the rotation generators (i.e. the angular momentum operators) are defined in such a way that they obey the convention of Varshalowich et al.[3]. The latter implies in particular the 'plus' sign on the right-hand sides of the corresponding commutators for the intrinsic-frame angular-momentum operators (it is sometimes forgotten that the often cited in this context the 'minus' sign is a matter of a representation - and of course no final result depends on that).

GENERALISED ROTORS: QUANTUM AND CLASSICAL HAMILTONIANS

The nuclear collective rotation can be often described using the rotor hamiltonian of the form

$$\hat{H} = \frac{\hat{I}_x^2}{2\mathcal{J}_x} + \frac{\hat{I}_y^2}{2\mathcal{J}_y} + \frac{\hat{I}_z^2}{2\mathcal{J}_z} \quad (1)$$

where $\hat{I}_x$, $\hat{I}_y$ and $\hat{I}_z$ denote the angular momentum operators in the body-fixed reference frame and the three constants, $\mathcal{J}_x$, $\mathcal{J}_y$ and $\mathcal{J}_z$ are interpreted as the effective moments of inertia. Such a hamiltonian is invariant with respect to the D_2 point group of symmetry and cannot be used to describe system of higher, e.g. tetrahedral symmetry.

In order to provide mathematical means needed to describe the motion of the higher-symmetry quantum objects we are going to introduce a basis of the spherical-tensor operators

defined by

$$\hat{T}_\mu^\lambda = \left(\left(\dots \left((\hat{\vec{I}} \otimes \hat{\vec{I}})^{\lambda_2=2} \otimes \hat{\vec{I}} \right)^{\lambda_3=3} \otimes \dots \otimes \hat{\vec{I}} \right)^{\lambda_{n-1}=n-1} \otimes \hat{\vec{I}} \right)_{\mu}^{\lambda_n=\lambda=n}, \quad (2)$$

with the help of the usual Clebsch-Gordan coupling and the corresponding generalized rotor hamiltonian as an expansion within the above basis:

$$\hat{H} = h_{00} T_0^0(2) + \sum_{\lambda=1}^{\infty} \left(h_{\lambda 0} \hat{T}_0^\lambda + \sum_{\mu=1}^{\lambda} \left(h_{\lambda \mu} \hat{T}_\mu^\lambda + (-1)^\mu h_{\lambda \mu}^* \hat{T}_{-\mu}^\lambda \right) \right), \quad (3)$$

where $T_0^0(2) = (\hat{\vec{I}} \otimes \hat{\vec{I}})_0^0$. Here, the symbol $\{h_{\lambda \mu}\}$ denotes the full set of the model parameters (constants); they however, may be some functions of the scalar expressions constructed out of $\vec{I}$. These parameters are defined in such a way that the resulting hamiltonian is hermitian. Within such a basis any rotor hamiltonian of any predefined point-group of symmetry can be easily constructed. Of course the standard rotor expression, Eq. (1), is a very particular case of that in Eq. (3), see also below.

To obtain a classical description of rotation that is compatible with that provided by a given quantum rotor hamiltonian we 'de-quantize' the latter using the *inverse* canonical quantization procedure:

$$\frac{1}{i\hbar} [\hat{f}, \hat{g}] \longrightarrow \{f, g\}, \quad (4)$$

where the symbols $[,]$ and $\{, \}$ refer to commutators and Poisson brackets, respectively. The three Euler-angles (formally: operators): $\hat{\alpha}$, $\hat{\beta}$, $\hat{\gamma}$ and the corresponding derivatives: $-i\frac{\partial}{\partial \alpha}$, $-i\frac{\partial}{\partial \beta}$ and $-i\frac{\partial}{\partial \gamma}$ should be replaced by the canonically conjugated variables α , β and γ , and the related momenta ℓ_α , ℓ_β , ℓ_γ , respectively.

Having de-quantized angular momenta we obtain a classical Hamiltonian - the classical analog of the original quantum Hamiltonian that was invariant with respect to a certain point-group of symmetry - and we can write down the Hamilton equations. In principle they form a system of six nonlinear differential equations. Using the fact that the angular momentum of an isolated object is conserved in the laboratory frame and selecting the $\mathcal{O}_z$ -axis of the latter to coincide with the direction of the corresponding angular-momentum vector, the first three of these equations can be reduced to the following form

$$\dot{\alpha} = \frac{\partial H}{\partial \ell_\alpha} = A(\beta, \gamma, |\vec{J}|), \quad \dot{\beta} = \frac{\partial H}{\partial \ell_\beta} = B(\beta, \gamma, |\vec{J}|), \quad \dot{\gamma} = \frac{\partial H}{\partial \ell_\gamma} = G(\beta, \gamma, |\vec{J}|), \quad (5)$$

valid in the body fixed coordinate frame. (Recall: here as always the Euler angles denote the relative angular position of the body-fixed *vs.* laboratory reference frame, and $|\vec{J}|$ is the

length of the angular momentum vector - a constant of the motion for a free rotor. Symbols A , B and G denote known functions of their arguments; their explicit form can only be defined after the symmetry group of interest has been chosen).

As one can see, the above equations do not depend on α . Consequently, solving the system of the second and third of the above equations for $\beta = \beta(t)$ and $\gamma = \gamma(t)$ and inserting the solutions into the equation containing $\dot{\alpha}$ allows to integrate directly the latter and obtain the full description of the relative motion of the two reference frames.

Concerning the other three Hamilton equations i.e. those containig $\dot{\ell}_\alpha$, $\dot{\ell}_\beta$ and $\dot{\ell}_\gamma$: by using the known relation between the laboratory and the body-fixed reference frames given by the solutions $\alpha(t)$, $\beta(t)$ and $\gamma(t)$ the components of $\vec{J}$ in the *body-fixed* reference frame can be expressed using only the length of the angular momentum vector (constant of the motion) and the angles β and γ . In other words: because of the conservation of the angular momentum in the laboratory reference frame, the motion of the angular momentum vector in the body-fixed reference frame can be obtained without effectively solving the latter three differential equations. Thus having found solutions only for β and γ one obtains the full description of the motion for the angular momentum vector in the body-fixed frame.

In the following we are going to be interested in particular in the equilibrium points (stable and unstable) associated with the rotational motion; these can be analyzed within the well established mathematical approach, cf. e.g. Ref.[4]. The behavior of the rotational motion in the vicinity of the equilibrium points ($\dot{\beta} = 0$, $\dot{\gamma} = 0$) allows to classify the trajectories of the classical rotor and relate them to the quantum spectrum. These relations will be described qualitatively in the rest of the paper.

The equilibrium points can be found from the conditions: $\dot{\beta} = 0$, $\dot{\gamma} = 0$.

In the following we are going to discuss the properties of the D_2 -symmetric rotor as a reference object to be compared with the higer symmetry (here: T_d) rotor.

D_2 -SYMMETRIC ROTOR

First let us consider the second order rotor with D_2 symmetry: its hamiltonian, Eq. (1), rewritten using the spherical tensor basis introduced in Eq. (2), takes the form

$$H_{\text{rot}} = h_{00} T_0^0(2) + h_{20} T_0^2 + h_{22} (T_2^2 + T_{-2}^2), \quad (6)$$

where $T_0^0(2) = (\vec{J} \otimes \vec{J})_0^0$; there is a one-to-one correspondence between the set of the three moments of inertia and the h -parameters in Eq. (6). In this case the Hamilton equations (5) read:

$$\dot{\alpha} = -\frac{J}{3} \left(\sqrt{6} h_{20} - 6 h_{22} \cos(2\gamma) \right); \quad (7)$$

$$\dot{\beta} = -2 J h_{22} \sin(\beta) \sin(2\gamma); \quad (8)$$

$$\dot{\gamma} = J \cos(\beta) \left(\sqrt{6} h_{20} - 2 h_{22} \cos(2\gamma) \right). \quad (9)$$

One can demonstrate, cf. Ref.[4], that the motion of the rotor in the neighborhood of equilibrium points found through $(\dot{\beta} = 0, \dot{\gamma} = 0)$, say (β_0, γ_0) , is determined by the eigenvalues of the Jacobi matrix, $\mathcal{M}$, defined as follows: suppose that the equations in question have the form

$$\dot{\beta} = B(\beta, \gamma, |\vec{J}|), \quad (10)$$

$$\dot{\gamma} = G(\beta, \gamma, |\vec{J}|). \quad (11)$$

Then the Jacobi matrix at the equilibrium point satisfies by definition

$$\mathcal{M} = \begin{bmatrix} \frac{\partial B}{\partial \beta}|_{(\beta_0, \gamma_0)}, & \frac{\partial B}{\partial \gamma}|_{(\beta_0, \gamma_0)} \\ \frac{\partial G}{\partial \beta}|_{(\beta_0, \gamma_0)}, & \frac{\partial G}{\partial \gamma}|_{(\beta_0, \gamma_0)} \end{bmatrix}. \quad (12)$$

It can be shown that in the discussed situation there are the following two possibilities. The solutions describing the motion of the angular momentum vector are periodic (closed trajectories) corresponding to the imaginary eigenvalues of the Jacobi matrix, or aperiodic ('hyperbolic') ones, corresponding to the real eigenvalues. In the latter case the equilibrium points are called saddles, in the former case - centers. The exact solutions for saddles are called separatrix trajectories.

For the illustration below we have selected the rigid-rotor moments of inertia corresponding to the quadrupole deformations $\beta_2 = 0.25$ and $\gamma = 30^\circ$, i.e. maximum triaxiality. After straightforward calculations we obtain the following pairs of equilibrium points for the discussed case: a. the first pair, $\{\beta, \gamma\} = \{\pi/2, 0\}$ and $\{\beta, \gamma\} = \{\pi/2, \pi\}$, and, b. the second pair $\{\beta, \gamma\} = \{\pi/2, \pi/2\}$ and $\{\beta, \gamma\} = \{\pi/2, 3\pi/2\}$.

The quantum spectra of the corresponding D_2 -symmetric hamiltonian have the following very well known behaviour: the states with the energies approaching the extremes in the

energy scale lie in pairs whose energies are very close to each other (cf. Fig. 1). With the energies approaching the middle of the scale the 'partnership' relation becomes weaker and weaker, corresponding energies lying further and further apart. One can show that among $(2I + 1)$ states in the spin I multiplet, the $2I$ states have a tendency to form the doublets of partners with very closely lying energies, one state staying always without a 'partner'.




FIG. 1: Energy surface of Eq. (13) with classical trajectories (left) and quantum spectrum (right) for D_2 -symmetric rotor. Hamiltonian coefficients are: $h_{00} = -0.016$, $h_{20} = 0.002$, $h_{22} = 0.001$, corresponding to the moments of inertia $J_x = 53.1$, $J_y = 67.4$ and $J_z = 45.9$ in $\hbar/\text{MeV}$ for the angular momentum $J = 10\hbar$ - i.e. a physical situation corresponding to a typical nucleus in Rare Earth nuclei.

The behaviour of the classical rotors can conveniently be discussed with the help of auxiliary surfaces on which the classical trajectories-solutions can be drawn. These surfaces are defined as follows. We introduce a body-fixed reference frame within which the motion of the angular momentum vector $\vec{J}(\alpha, \beta, \gamma)$ can be treated as already known in terms of the functions $\{\alpha(t), \beta(t), \gamma(t)\}$. To each orientation of this vector there corresponds rotor's energy $E(\alpha, \beta, \gamma)$; the surface composed of all points $\vec{\xi}$ below

$$\vec{\xi} = E(\alpha, \beta, \gamma) \frac{\vec{J}(\alpha, \beta, \gamma)}{|\vec{J}(\alpha, \beta, \gamma)|} \quad (13)$$

will serve as our reference surface (cf. Figs. 1 and 2). In other words: this surface is composed of all the points corresponding to all the solutions of the motion associated to a given length of the vector $\vec{J}$; consequently all trajectories - solutions to the Hamilton equations with $|\vec{J}|$ -given must lie on such a surface.

Comparing the quantum spectra for the D_2 -symmetric Hamiltonian with the classical solutions whose energies are exactly equal to those of the corresponding quantum case, one can observe several correlations. First of all, the energy associated with the separatrix is close to the energy of the partnerless state and in this sense the two objects can be associated. Secondly, states lying above and below the separatrix correspond to the closed trajectories surrounding centers with maximal and minimal energy. One can observe higher density of states when energy of the system is approaching the energy associated with the separatrix.

T_d -SYMMETRIC ROTOR

The simplest hamiltonian of the generalized rotor with tetrahedral symmetry has the form:

$$\hat{H}_t = \underbrace{h_{00} T_0^0(2)}_{H_0} + \underbrace{h_{32} (T_2^3 - T_{-2}^3)}_{H_3}, \quad (14)$$

so that to guarantee the hermiticity of the rotor hamiltonian the h_{32} coefficients must be purely imaginary. In the classical limit we have

$$T_2^3 - T_{-2}^3 \rightarrow i I_x I_y I_z. \quad (15)$$

This structure implies the time-odd character of the latter term since the i -factors disappear and the product of the three I -operators changes sign under time reversal.

The latter feature is worth examining because it implies an interesting property of the whole hamiltonian in Eq. (14). Since $[\hat{H}_0, \hat{H}_3] = 0$, it follows that the eigenvalues of $\hat{H}_t$, satisfy $E_t = E_0 + E_3$, where E_0 and E_3 are the eigenvalues of $\hat{H}_0$ and $\hat{H}_3$, respectively. We may write

$$\hat{H}_t \Psi_t = E_t \Psi_t \rightarrow (\hat{H}_0 + \hat{H}_3) \Psi_t = E_t \Psi_t \quad (16)$$

and at the same time

$$\hat{\mathcal{T}} \hat{H}_t \hat{\mathcal{T}}^{-1} = \hat{H}_0 - \hat{H}_3 \rightarrow (\hat{H}_0 - \hat{H}_3) \Psi_t^* = E_t \Psi_t^* \quad (17)$$

and comparing the relations (16) and (17) we find

$$\hat{H}_3 \Psi_t = +(E_t - E_0) \Psi_t \quad \leftrightarrow \quad \hat{H}_3 \Psi_t^* = -(E_t - E_0) \Psi_t^* \quad (18)$$

i.e. the spectrum of the $\hat{H}_3$ term in the hamiltonian is symmetric with respect to zero. Assuming that the energy of the rotational motion is 'purely cinetic' we must select the paramters of the full hamiltonian $\hat{H}_t$ in such a way that the quadtratic (scalar) term dominates so that E_t remains positive.

In principle the nuclear rotor hamiltonian is a sum of at least two terms one of them representing the intrinsic structure of the nucleus and the other one the coherent rotational motion; only the sum of the two has a priori definite transformation properties with respect to the symmetry operations (so has the total wave function corresponding to such a hamiltonian). Following this view point one needs to assume that h_{32} coefficient is a function of group symmetry invariants e.g. J^2 and y^{int} the latter describing the internal-coordinate structure of the rotating body. This function is odd under the time reversal operation like the third-order term in Hamiltonian and the product of them would allow to conserve the time reversal symmetry. Such an interpretation would modify the approach to the highly symmeric rotors, here, however, we are not going to discuss this more complete model of the nuclear rotor.




FIG. 2: Energy surface with classical trajectories and quantum spectrum for T_d -symmetric rotor. Hamiltonian coefficients are: $h_{00} = -0.016$, $h_{32} = i0.003003$, $J = 16$.

The Hamilton equations for the Hamiltonian (16) are:

$$\dot{\alpha} = -i 2\sqrt{3} J^2 h_{32} \cos(\beta) \sin(2\gamma); \quad (19)$$

$$\dot{\beta} = -i \sqrt{3} J^2 h_{32} \cos(2\beta) \sin(2\gamma); \quad (20)$$

$$\dot{\gamma} = \frac{i\sqrt{3}}{2} J^2 h_{32} \sin(2\gamma)(1 + 3 \cos(2\beta)); \quad (21)$$

they are real since the h_{32} -coefficient is purely imaginary.

Calculations show that there are eight equilibrium points for this system corresponding to all possible combinations of $\beta = \arccos(1/3)$, $\pi - \arccos(1/3)$, $\gamma = \pi/4, 3\pi/4, 5\pi/4, 7\pi/4$ and the four points related to $\beta = \pi/2$ and $\gamma = 0, \pi/2, \pi, 3\pi/2$, classified as saddles.

Analyzing the classical-motion picture, Fig. 2, and the quantum spectrum one can observe an increase of the density of states near the energy associated with the separatrix. Another feature of the quantum spectrum, explained by Eq. (19), is that it is symmetric (in the sense of Eq. (19)) with respect to the state with the energy equal to that of the separatrix-solution. The symmetry in question corresponds in fact to an exchange of the roles of the two (out of three) one-dimensional irreducible representations of the T_d -group. In particular one may note that at an even spin value, $I=16$ in the illustrated case, all the states below the separatrix form multiplets with the energy of the three-dimensional irrep *above* the energies of the one-dimensional ones - this tendency is exactly opposite above the separatrix.

At this stage we have no further comments about possible analogies between the features of classical trajectories and e.g. triple degeneracies of quantum states corresponding to the 3-dimensional irreducible representation of the T_d -group.

SUMMARY AND CONCLUSIONS

We have constructed a mathematical scheme allowing to draw certain parallels between the quantum behaviour of high-symmetry rotors and the solutions of the implied classical equations of the motion for the angular-momentum vector in the intrinsic (body-fixed) reference frame. The procedure is analogous to those used in certain studies in molecular physics. Unlike molecular physics applications where often the semi-classical or classical limits are of interest ('angular momentum $\rightarrow \infty$ '), the nuclear physics applications must focus on the very low-spin limits, since the angular-momentum alignment phenomena are expected to destroy the delicate balance leading to such high symmetries. Moreover, the

total symmetries of the systems are very different in the two types of objects (for instance, the symmetry with respect to exchange of positions of nuclei/atoms in the molecules have no nuclear physics analogs). This latter aspect is in general not just a trivial 'tiny' complication - the full group-structure and thus the related irreducible representations (even their number), the tensor character of the implied wave functions and their symmetry/antisymmetry properties are influenced. This can be viewed as a challenge for the nuclear structure physics whose observables and physics-cases seem to be significantly different in many respects from those in the the traditional molecular physics applications.

- [1] J. Dudek, A. Gózdź, N. Schunck and M. Miśkiewicz, Phys. Rev. Lett. **88**, 252502 (2002);
 J. Dudek, A. Gózdź and D. Rosły, Acta Phys. Polonica. **B32** 2625 (2001);
 A. Gózdź, J. Dudek and M. Miśkiewicz, Acta Phys. Polon. **B34** 2123 (2003);
 J. Dudek, A. Gózdź, N. Schunck, Acta Phys. Polonica. **B34** 2491 (2003).
- [2] A. Barut, R. Rączka; *Theory of Group Representations and Applications* (PWN, Warszawa, 1977).
- [3] D. Varshalovich, A. Moskalev and V. Khershonskii; *Quantum Theory of Angular Momentum* (WSP, Singapore, 1988) Ch. 4.2, p.75.
- [4] J. Hubbard, B. West; *Differential Equations: A Dynamical Systems Approach* (Springer-Verlag, 1990), Ch. 8.1, p.137.